STATE OF THE ART

2005 emergency cardiovascular care guidelines

Michael Shuster, MD

Introduction

The 2005 Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care (referred to here as the "2005 ECC guidelines") recommend major changes to basic resuscitation practices and significant changes to advanced care.¹ This article highlights some of those changes and gives a brief overview of the guidelines creation process.

Creating the 2005 ECC guidelines

The comprehensive guidelines creation process followed in 2005 has evolved significantly beyond the process used to create previous guidelines.

In 1992, the Heart and Stroke Foundation of Canada (HSFC) participated in the American Heart Association's (AHA) guideline process and then adapted the AHA guidelines for Canada.

In 2000, the HSFC partnered with the AHA and other members of the International Liaison Committee on Resuscitation (ILCOR)* to develop the first international ECC guidelines. Consensus on some issues was difficult or impossible to reach because of differences in resources and medical practice in different parts of the world. Even after the guidelines were agreed upon, each of the international participants then altered the guidelines for their home countries.

In 2005, rather than try to reach an international consen-

sus on guidelines, ILCOR agreed to develop an international consensus on science and also on treatment recommendations where possible, but to leave the establishment of guidelines to each of its members. The HSFC decided that, rather than "Canadianize" the AHA guidelines as it had done before 2000, HSFC would work with the AHA to develop a single set of guidelines that would be acceptable in both Canada and the United States.

This 2005 guideline revision is the result of a process that began 5 years ago, immediately following the completion of the 2000 guidelines. The process has involved thousands of volunteer hours by hundreds of resuscitation experts and researchers from around the world, who reviewed 276 topics and completed more than 400 worksheets. In January 2005, 380 experts from around the world met in Dallas under the auspices of ILCOR to examine the evidence on resuscitation and to develop a consensus on science and on treatment recommendations. This international consensus forms the basis for the new guidelines. The guidelines themselves are a result of a collaboration between the AHA and the HSFC.

Previous guidelines have been criticized for reflecting the influence of pharmaceutical companies and other special interests. For the 2005 guidelines, ILCOR and the AHA went to great lengths to ensure full disclosure of potential conflicts and to manage conflict of interest when it existed. Details of the conflict of interest policy have been published along with the consensus document.²

Why create new ECC guidelines

The purpose of guidelines is to encourage rational treatment decisions with a view to reducing preventable mor-

Department of Emergency Medicine, Mineral Springs Hospital, Banff, Alta.

Received: Nov. 13, 2005; accepted: Nov. 21, 2005

Can J Emerg Med 2006;8(1):37-42

^{*}The Heart and Stroke Foundation of Canada, European Resuscitation Council, the American Heart Association, the Resuscitation Council of Southern Africa, the Australia and New Zealand Resuscitation Council, and the Inter-American Heart Foundation.

bidity and mortality. These resuscitation guidelines provide advice for best practice based on the best available evidence, and on expert consensus when sufficient evidence is not available.

The guidelines are intended to assist practitioners who perform resuscitation sporadically, or even regularly, to take a rational approach to the procedure. The guidelines serve as a substitute for individual research, as it is unrealistic for most care providers to evaluate all of the existing evidence around every resuscitation situation they encounter. These guidelines are not intended to lock physicians into a particular care regimen, as "rules" or "standards" would. Since there is always lag time between the publication of evidence and the incorporation of that evidence into guidelines, the expert who is familiar with all of the latest evidence and who is able to critically review that evidence may well choose to deviate from guidelines that have not yet taken new research into account.

Because of the need for guidelines to make recommendations even when the evidence is weak (i.e., caregivers cannot opt not to care for patients because there is no definitive evidence of therapeutic benefit) some of the recommendations may be controversial. While caregivers do not want to rush into changes that may at best be unhelpful, or at worst cause harm, they cannot be paralyzed into inaction by a lack of definitive evidence in a condition where mortality is 95% or higher and definitive evidence is extremely difficult to develop.

Major ECC guidelines changes

Table 1 illustrates the major changes in the 2005 ECC guidelines. Few of the changes are based on clear-cut evidence from large randomized controlled trials. While a great deal of research has been done on resuscitation in the past 5 years, little is certain beyond the knowledge that early cardiopulmonary resuscitation (CPR) and early defibrillation are important determinants of survival.

The story behind ILCOR's development of an updated algorithm for CPR provides an excellent demonstration of how the committee dealt with gaps in the science of resuscitation.

There are no human studies showing that the new CPR algorithm produces better outcomes than the previous CPR algorithm. However, there is a great deal of circumstantial evidence to suggest that it should.

Compression:ventilation ratio: In order to be successful, CPR must develop and maintain adequate myocardial perfusion pressure.^{3–5} The ILCOR committee concluded that the 2000 algorithm is not optimal at achieving this aim. Two recent studies show that interruptions in chest compressions are frequent and, for a substantial proportion of arrest time, no compressions are provided.^{6.7}

Coronary perfusion pressure is lost each time CPR is paused, and it then takes 3 to 5 compressions to develop adequate perfusion pressure when CPR is resumed.⁸⁻¹⁰ While it may seem that pausing for a couple of seconds to ventilate doesn't greatly affect "no perfusion" time, studies show that it takes a team of 2 rescuers ~8 seconds, or a single rescuer ~15 seconds, to perform a set of ventilations.^{11,12} At a ratio of 15:2, and a rate of 100 compressions per minute, a single rescuer therefore provides only about 45 compressions per minute — of which only 33 are effective — along with 5 ventilations per minute, while 2 rescuers deliver 44 effective compressions per minute (60 actual compressions) along with 6 ventilations.

One human study and multiple animal studies clearly show that higher compression rates (>80) without interruption are more effective in non-asphyxial arrest than compressions with many or prolonged interruptions.^{9,13–21} In an analysis of ventricular fibrillation (VF) waveform, the interruption of CPR was associated with a lower probability of conversion of VF to another rhythm.²² In both manikin and animal studies, coronary and cerebral perfusion pressure increases as chest compression rates increase.^{5,23,24}

Compression:ventilation ratios of 100:2 and 50:2 have received limited study with mixed results.^{25,26} Rescuer fatigue at high continuous compression rates may be a limiting factor.²⁷ In the single animal study where it was tried, a ratio of 30:2 was associated with a significantly shorter time to return of spontaneous circulation, and greater systemic and cerebral oxygenation as compared with continuous chest compressions.²⁸ Finally, a theoretical analysis of various compression:ventilation ratios suggests 30:2 would provide the best blood flow and oxygen delivery.²⁹

Therefore, with a variety of circumstantial and low-level evidence, and with a view toward simplifying teaching, promoting skills retention, increasing the number of compressions and decreasing the interruptions to compressions, the consensus of ILCOR experts was that a change to a ratio of 30:2 was warranted.

Single versus stacked shocks: Another change to the guidelines that will have a positive effect on decreasing "no perfusion" time is the elimination of "stacked shocks," which have been a part of the guidelines for many years. The practice was introduced because of evidence that trans-thoracic impedance decreased with each successive shock and that lower energy shocks were not effective in patients with high impedance.^{30,31}

The new guidelines replace stacked shocks with a single (preferably biphasic) shock. With a first-shock efficacy of around 90% for biphasic defibrillation, stacked shocks provide little incremental value and unduly delay compressions.³²⁻⁴⁰ This change will have a major impact on "no perfusion" time for all defibrillator uses, but especially when an automatic external defibrillator (AED) is used because programmed re-analysis and shock may take up to 55% of resuscitation time.³⁶ Yu and colleagues demonstrated experimentally that there is a dramatic increase in the predicted return of spontaneous circulation when defibrillation is not preceded by a long pause in compressions.⁴¹

Recommended shock energy for biphasic defibrillation

Table 1. Major differences between the 2005	Guidelines for Cardiopulmonary Resuscitation and Emergency
Cardiovascular Care and the 2000 guidelines	

	Major guideline differences	
Item	New 2005 guideline	Old 2000 guideline
Compression:Ventilation ratio – Adult, with unprotected airway (compression rate 100/min)	30:2	15:2
Compression:Ventilation ratio – Pediatric, with unprotected airway (compression rate 100/min)	30:2 (lay rescuers and single health care provider) or15:2 (two health care providers)	15:2 (lay rescuers and single health care provider) or5:1 (two health care providers)
Ventilation rate (with protected airway)	Max 8–10 per minute	10–12 per minute
Duration of CPR between shocks	2 minutes	1 minute
Shocks per defibrillation attempt	Single shock, preferably biphasic	3 escalating energy "stacked" shocks
CPR quality	 limit interruptions ensure adequate depth of compression allow full chest recoil don't hyperventilate 	Not emphasized
Pulse check post-shock	No Resume CPR immediately post-shock	Yes
Rhythm check post-shock	No Resume CPR immediately post-shock	Yes
Advanced airways	LMA, Combitube™ and tracheal tube are equally acceptable	LMA and Combitube™ recommended over bag-mask when tracheal tube not available
Vasopressors	Give after one unsuccessful shock followed by 2 minutes of CPR if still no perfusing rhythm	Give after series of 3 unsuccessful shocks
Anti-arrhythmics	"Consider" after 2nd unsuccessful shock followed by 2 minutes of CPR if still in VF	"Consider" after unsuccessful stacked-shock series and second unsuccessful shock
Atropine for asystole	"Consider" for brady-asystolic arrest	"Recommended"
Fibrinolysis during cardiac arrest	Consider if suspected pulmonary embolism or when acute thrombotic etiology suspected in initially failed resuscitation	N/A
Treatment of stable tachycardia out-of-hospital	Limit prehospital treatment of stable narrow complex tachycardia to vagal manoeuvres or adenosine. Defer treatment of stable wide- complex tachycardia until patient is in hospital.	Complex algorithm
Post cardiac arrest care	Cool unconscious adult patients with spon- taneous circulation to 32–34°C for 12–24 hr ("recommended" post-VF out-of-hospital arrest, "consider" post non-VF out-of-hospital arrest or any in-hospital arrest)	N/A

CPR = cardiopulmonary resuscitation; LMA = laryngeal mask airway; Combitube (Tyco/Kendall, Mansfield, Mass.) = a double lumen tube, one lumen resembling a tracheal tube, the other an esophageal obturator type tube with a distal blocked end and perforations at the pharyngeal level; VF = ventricular fibrillation

is 200 J if the device-specific energy setting is not known. Because devices use different biphasic waveforms, devicespecific shock energy will vary. High first shock efficacy has been demonstrated for energy levels ranging from 120 J to 200 J depending on the specific device used. Subsequent shocks can safely and effectively be given at 150–360 J.

If a monophasic defibrillator is used, the new recommendation is to deliver a single shock of 360 J. Under the previous guidelines, the first shock was performed at 200 J because of a single study that showed a higher rate of postshock atrioventricular heart block using 320 J.⁴² The postshock heart block that was demonstrated at this higher setting was transitory, and the experts now concur that a single energy level setting of 360 J for all monophasic shocks is reasonable.

Post-shock rhythm analysis and pulse checks: The 2005 guidelines recommend that CPR resume immediately following defibrillation, thus eliminating the intervening steps of the post-shock rhythm analysis and pulse check that had been previously recommended. Not only do these post-shock interventions result in a substantial "no perfusion" time, but studies indicate that pulse checks are unreliable, and analyses of electrical activity and of perfusion pressures suggest that delaying compressions post-shock is detrimental to VF conversion and to survival.^{22,39,41,43-45} Furthermore, there is evidence that chest compressions do not adversely affect a patient with an organized rhythm and a minimally perfusing rhythm.⁴⁶ Taken together, the evidence supports the conclusion that there is more benefit than risk in the immediate resumption of CPR following defibrillation.

Drug administration: Recommendations regarding vasopressors and anti-arrhythmic drugs were both subjects of controversy following the 2000 guidelines. Since that time, little new evidence has emerged. There is still no placebocontrolled trial of vasopressors in cardiac arrest, and new studies comparing vasopressin to epinephrine have not cast any light on the issue. A meta-analysis of 5 randomized controlled trials comparing vasopressin to epinephrine shows no difference between the 2 drugs for any outcome.47 Nor are there any placebo-controlled trials of antiarrhythmics to guide therapy. A study that compared amiodarone to lidocaine, while favouring amiodarone to lidocaine for survival to hospital admission, was not powered to examine survival to hospital discharge. Given the lack of new evidence, the 2005 guidelines have not changed with respect to vasopressors (which are recommended) and anti-arrhythmics (which should be considered). However, the timing and frequency of administration does change substantially.

The 2000 guidelines recommended the following sequence: shock, rhythm analysis, pulse check, vasopressor. The new guidelines recommend the immediate resumption of CPR after the first shock, with the first post-shock rhythm analysis taking place following 2 minutes of CPR. Vasopressor is administered after this analysis (i.e., 2 minutes later in the course of the arrest than the old guidelines recommended) — but only if administration will not delay a second shock. If giving the vasopressor will delay the shock, it should be administered immediately after the second shock. Because there is no evidence that vasopressors improve survival, the new guidelines give priority to uninterrupted CPR at the expense of earlier administration of vasopressor. Similarly, because there is no evidence that anti-arrhythmics improve survival, the experts felt that consideration of anti-arrhythmic administration should occur only after a second shock fails to convert the rhythm from ventricular fibrillation (i.e., 3 minutes later in the course of the arrest than the 2000 guidelines recommended).

Conclusion

The 2005 ECC guidelines are a work in progress. The last 30 years have seen the development of a much more evidence-based approach to medical science, an expansion of our knowledge of resuscitation and related issues, and far broader involvement of the international scientific community in sharing expertise, ideas and creativity. There remain glaring gaps in our knowledge of resuscitation science. It is hoped that this iteration of the guidelines, along with the accompanying controversies, will, as previous iterations have, highlight new avenues for exploration, raise questions for which we do not yet have answers and inspire research that will lead to improved future guidelines and ever better outcomes for our patients.

Competing interests: Michael Shuster chairs the Heart and Stroke Foundation of Canada's Policy Advisory Committee on Resuscitation. He has been involved with Emergency Cardiovascular Care guidelines review since 1992, actively participating in the literature evaluation and consensus meetings that led to the 2005 Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care.

Key words: guidelines, emergency cardiovascular care; resuscitation

References

1. Emergency Cardiovascular Care Committee and Subcommittees of the American Heart Association. 2005 American Heart Asso-

ciation Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Circulation 2005;112:IV-1–IV-211.

- International Liaison Committee on Resuscitation. 2005 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. Circulation 2005;112(suppl III):III-1–III-136.
- Paradis NA, Martin GB, Rivers EP, et al. Coronary perfusion pressure and the return of spontaneous circulation in human cardiopulmonary resuscitation. JAMA 1990;263:1106-13.
- Kern KB, Ewy GA, Voorhees WD, et al. Myocardial perfusion pressure: a predictor of 24-hour survival during prolonged cardiac arrest in dogs. Resuscitation 1988;16:241-50.
- Halperin HR, Tsitlik JE, Guerci AD, et al. Determinants of blood flow to vital organs during cardiopulmonary resuscitation in dogs. Circulation 1986;73:539-50.
- Abella BS, Alvarado JP, Myklebust H, et al. Quality of cardiopulmonary resuscitation during in-hospital cardiac arrest. JAMA 2005;293:305-10.
- Wik L, Kramer-Johansen J, Myklebust H, et al. Quality of cardiopulmonary resuscitation during out-of-hospital cardiac arrest. JAMA 2005;293:299-304.
- Sanders AB, Ogle M, Ewy GA. Coronary perfusion pressure during cardiopulmonary resuscitation. Am J Emerg Med 1985; 3:11-4.
- Berg RA, Sanders AB, Kern KB, et al. Adverse hemodynamic effects of interrupting chest compressions for rescue breathing during cardiopulmonary resuscitation for ventricular fibrillation cardiac arrest. Circulation 2001;104:2465-70.
- Sanders ABea. Importance of the duration of inadequate coronary perfusion pressure on resuscitation from cardiac arrest. J Am Coll Cardiol 1985;6:113-8.
- Assar D, Chamberlain D, Colquhoun M, et al. Randomised controlled trials of staged teaching for basic life support, 1: skill acquisition at bronze stage. Resuscitation 2000;45:7-15.
- 12. Heidenreich JW, Higdon TA, Kern KB, et al. Single-rescuer cardiopulmonary resuscitation: "two quick breaths" an oxymoron. Resuscitation 2004;62:283-9.
- 13. Swenson RD, Weaver WD, Niskanen RA, et al. Hemodynamics in humans during conventional and experimental methods of cardiopulmonary resuscitation. Circulation 1988;78:630-9.
- 14. Kern KB, Sanders AB, Raife J, et al. A study of chest compression rates during cardiopulmonary resuscitation in humans: the importance of rate-directed chest compressions. Arch Intern Med 1992;152:145-9.
- Berg RA, Cobb LA, Doherty A, et al. Chest compressions and basic life support-defibrillation. Ann Emerg Med 2001;37(4 suppl):S26-35.
- 16. Berg RA, Hilwig RW, Kern KB, et al. Automated external defibrillation versus manual defibrillation for prolonged ventricu-

lar fibrillation: lethal delays of chest compressions before and after countershocks. Ann Emerg Med 2003;42:458-67.

- 17. Kern KB, Hilwig RW, Berg RA, et al. Importance of continuous chest compressions during cardiopulmonary resuscitation: improved outcome during a simulated single lay-rescuer scenario. Circulation 2002;105:645-9.
- 18. Ewy GA. Cardiocerebral resuscitation: the new cardiopulmonary resuscitation. Circulation 2005;111:2134-42.
- 19. Berg RA, Kern KB, Hilwig RW, et al. Assisted ventilation does not improve outcome in a porcine model of single-rescuer bystander cardiopulmonary resuscitation. Circulation 1997;95: 1635-41.
- Berg RA, Kern KB, Hilwig RW, et al. Assisted ventilation during "bystander" CPR in a swine acute myocardial infarction model does not improve outcome. Circulation 1997;96:4364-71.
- Berg RA, Hilwig RW, Kern KB, et al. "Bystander" chest compressions and assisted ventilation independently improve outcome from piglet asphyxial pulseless "cardiac arrest." Circulation 2000;101:1743-8.
- 22. Effestol T, Sunde K, Steen PA. Effects of interrupting precordial compressions on the calculated probability of defibrillation success during out-of-hospital cardiac arrest. Circulation 2002;105: 2270-3.
- 23. Handley AJ, Handley JA. The relationship between rate of chest compression and compression:relaxation ratio. Resuscitation 1995;30:237-41.
- 24. Feneley MP, Maier GW, Kern KB, et al. Influence of compression rate on initial success of resuscitation and 24 hour survival after prolonged manual cardiopulmonary resuscitation in dogs. Circulation 1988;77:240-50.
- 25. Sanders AB, Kern KB, Berg RA, et al. Survival and neurologic outcome after cardiopulmonary resuscitation with four different chest compression-ventilation ratios. Ann Emerg Med 2002;40: 553-62.
- Dorph E, Wik L, Stromme TA, et al. Quality of CPR with three different ventilation:compression ratios. Resuscitation 2003;58: 193-201.
- 27. Greingor JL. Quality of cardiac massage with ratio compression-ventilation 5/1 and 15/2. Resuscitation 2002;55:263-7.
- Dorph E, Wik L, Stromme TA, et al. Oxygen delivery and return of spontaneous circulation with ventilation:compression ratio 2:30 versus chest compressions only CPR in pigs. Resuscitation 2004;60:309-18.
- 29. Babbs CF, Kern KB. Optimum compression to ventilation ratios in CPR under realistic, practical conditions: a physiological and mathematical analysis. Resuscitation 2002;54:147-57.
- Dahl CF, Ewy GA, Ewy MD, et al. Transthoracic impedance to direct current discharge: effect of repeated countershocks. Med Instrum 1976;10:151-4.
- 31. Geddes LA, Tacker WA, Cabler P, et al. The decrease in

transthoracic impedance during successive ventricular defibrillation trials. Med Instrum 1975;9:179-80.

- 32. Bain AC, Swerdlow CD, Love CJ, et al. Multicenter study of principles-based waveforms for external defibrillation. Ann Emerg Med 2001;37:5-12.
- 33. Mittal S, Ayati S, Stein KM, et al. Comparison of a novel rectilinear biphasic waveform with a damped sine wave monophasic waveform for transthoracic ventricular defibrillation. ZOLL Investigators. J Am Coll Cardiol 1999;34:1595-601.
- Poole JE, White RD, Kanz KG, et al. Low-energy impedancecompensating biphasic waveforms terminate ventricular fibrillation at high rates in victims of out-of-hospital cardiac arrest. LIFE Investigators. J Cardiovasc Electrophysiol 1997;8:1373-85.
- 35. Schneider T, Martens PR, Paschen H, et al. Multicenter, randomized, controlled trial of 150-J biphasic shocks compared with 200- to 360-J monophasic shocks in the resuscitation of out-of-hospital cardiac arrest victims. Circulation 2000;102: 1780-7.
- van Alem AP, Sanou BT, Koster RW. Interruption of cardiopulmonary resuscitation with the use of the automated external defibrillator in out-of-hospital cardiac arrest. Ann Emerg Med 2003;42:449-57.
- Martens PR, Russell JK, Wolcke B, et al. Optimal Response to Cardiac Arrest study: defibrillation waveform effects. Resuscitation 2001;49:233-43.
- Carpenter J, Rea TD, Murray JA, et al. Defibrillation waveform and post-shock rhythm in out-of-hospital ventricular fibrillation cardiac arrest. Resuscitation 2003;59:189-96.
- Gliner BE, White RD. Electrocardiographic evaluation of defibrillation shocks delivered to out-of-hospital sudden cardiac arrest patients. Resuscitation 1999;41:133-44.
- 40. White RD, Blackwell TH, Russell JK, et al. Transthoracic im-

pedance does not affect defibrillation, resuscitation or survival in patients with out-of-hospital cardiac arrest treated with a nonescalating biphasic waveform defibrillator. Resuscitation 2005; 64:63-9.

- 41. Yu T, Weil MH, Tang W, et al. Adverse outcomes of interrupted precordial compression during automated defibrillation. Circulation 2002;106:368-72.
- 42. Weaver WD, Cobb LA, Copass MK, et al. Ventricular defibrillation: a comparative trial using 175-J and 320-J shocks. N Engl J Med 1982;307:1101-6.
- 43. Eberle B, Dick WF, Schneider T, et al. Checking the carotid pulse check: diagnostic accuracy of first responders in patients with and without a pulse. Resuscitation 1996;33:107-16.
- 44. Moule P. Checking the carotid pulse: diagnostic accuracy in students of the healthcare professions. Resuscitation 2000;44: 195-201.
- Eftestol T, Wik L, Sunde K, Steen PA. Effects of cardiopulmonary resuscitation on predictors of ventricular fibrillation defibrillation success during out-of-hospital cardiac arrest. Circulation 2004: 01.CIR.0000133323.15565.75.
- 46. Hess E, White R. Ventricular fibrillation is not provoked by chest compression during post-shock organized rhythms in outof-hospital cardiac arrest. Resuscitation 2005;66:7-11.
- 47. Aung K, Htay T. Vasopressin for cardiac arrest: a systematic review and meta-analysis. Arch Intern Med 2005;165:17-24.

Correspondence to: Dr. Michael Shuster, Mineral Springs Hospital, Box 1050, Banff AB T1L 1H7; mtn1ms@poky.ca

The complete 2005 ECC guidelines are available through your provincial Heart and Stroke Foundation or online at www .heartandstroke.ca/